

Equations

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Diagram illustrating the components of the equation $5x + 7 = \sqrt{2}$:

- Coefficient** (green) points to 5 .
- Variable** (red) points to x .
- Constant** (yellow) points to 7 and $\sqrt{2}$.

The equation is structured as follows:

$$\underbrace{5x + 7}_{\text{Expression}} = \underbrace{\sqrt{2}}_{\text{Expression}}$$

The entire equation is labeled as an **Equation**.

Terms: $5x$, 7 , $\sqrt{2}$

MATHS

CA FOUNDATION

CODE : SS12

Equations



Shivani
Sharma

The
variable

Equality

$$2x - 3 = 7$$

Equation

$$2x - 3 = \text{LHS}$$

$$7 = \text{RHS}$$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Definition

- Equation is defined to be a mathematical statement of equality.

Example:

- $5x = 25$
- $2x - 3 = 9$
- $2y + \frac{5}{2} = \frac{37}{2}$
- $6z + 10 = -2$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Solution / Root of the equation

- The value of the variable which satisfies an equation is called the "solution" of the equation, or the "root" of the equation.

MATHS

CA FOUNDATION

CODE : SS12

CONDITIONAL EQUATION

If the equality is true for certain value of the variable involved , the equation is often called a conditional equation and equality sign ' $=$ ' is used .

For Example: $\frac{x+2}{3} + \frac{x+3}{2} = 3$ holds true only for $x=1$.

So it is a conditional.



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12

IDENTITY

If the equality is true for all values of the variable involved , the equation is called an identity .

EXAMPLE :

$$\frac{x+2}{3} + \frac{x+3}{2} = \frac{5x+13}{6}$$

is an identity since it holds for all values of the variable x .



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12

Degree



Shivani
Sharma

- The highest power of variable in a given equation is called degree.
- $8x + 17(x-3) = 4(4x-9) + 12$ is a **Linear equation**.
- $3x^2 + 5x + 6 = 0$ is a **Quadratic equation**.
- $3x^3 + 5x^2 + 6x + 1 = 0$ is a **Cubic equation**.

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

- **Simultaneous linear Equations.**
- Two or more linear equations involving two or more variables are called simultaneous linear equations.
- $x + 2y = 1$, $2x + 3y = 2$ are jointly called Simultaneous equations.

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Simple Equation

- Equation of one degree and having one unknown variable is simple equation.
- Simple equation has only one root.

General form

$$ax + b = 0$$

where ,a is coefficient of x , $a \neq 0$

b is constant

MATHS

EXERCISE (A)

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Que.1 The equation $-7x + 1 = 5 - 3x$ will be satisfied for x equal to:

- a. 2
- b. -1
- c. 1
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.2 The root of the equation $\frac{x+4}{4} + \frac{x-5}{3} = 11$ is

- a. 20
- b. 10
- c. 2
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.3 Pick up the correct value of x for $\frac{x}{30} = \frac{2}{45}$

- a. $x = 5$
- b. $x = 7$
- c. $x = 1\frac{1}{3}$
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.4 The solution of the equation $\frac{x + 24}{5} = 4 + \frac{x}{4}$

- a. 6
- b. 10
- c. 16
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.5 8 is the solution of the equation

a. $\frac{x+4}{4} + \frac{x-5}{3} = 11$

b. $\frac{x+4}{2} + \frac{x+10}{9} = 8$

c. $\frac{x+24}{5} = 4 + \frac{x}{4}$

d. $\frac{x-15}{10} + \frac{x+5}{5} = 4$

MATHS

CA FOUNDATION

CODE : SS12

Que.6

The value of y that satisfies the equation $\frac{y+11}{6} - \frac{y+1}{9} = \frac{y+7}{4}$ is

- a. -1
- b. 7
- c. 1
- d. -1/7



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.7 The solution of the equation $(p+2)(p-3) + (p+3)(p-4) = p(2p-5)$ is

- a. 6
- b. 7
- c. 5
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.8 The equation $\frac{12x+1}{4} = \frac{15x-1}{5} + \frac{2x-5}{3x-1}$ is true for

- a. $x = 1$
- b. $x = 2$
- c. $x = 5$
- d. $x = 7$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.9 Pick up the correct value x for which $\frac{x}{0.5} - \frac{1}{0.05} + \frac{x}{0.005} - \frac{1}{0.0005} = 0$

- a. $x = 0$
- b. $x = 1$
- c. $x = 10$
- d. None of these

MATHS

EXERCISE (B)

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Que.1 The sum of two numbers is 52 and their difference is 2. The numbers are

- a. 17 and 15
- b. 12 and 10
- c. 27 and 25
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.2 The diagonal of a rectangle is 5 cm and one of its sides is 4 cm.
Its area is

- a. 20 sq. cm
- b. 12 sq. cm
- c. 10 sq. cm
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.3 Divide 56 into two parts such that three times the first part exceeds one third of the second by 48. The parts are.

- a. (20, 36)
- b. (25, 31)
- c. (24, 32)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.4 The sum of the digits of a two digit number is 10. If 18 be subtracted from it the digits in the resulting number will be equal. The number is

- a. 37
- b. 73
- c. 75
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.5 The fourth part of a number exceeds the sixth part by 4. The number is

- a. 84
- b. 44
- c. 48
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.6 Ten years ago the age of a father was four times of his son. Ten years hence the age of the father will be twice that of his son. The present ages of the father and the son are.

- a. (50, 20)
- b. (60, 20)
- c. (55, 25)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.7 The product of two numbers is 3200 and the quotient when the larger number is divided by the smaller is 2. The numbers are

- a. (16, 200)
- b. (160, 20)
- c. (60, 30)
- d. (80, 40)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.8 The denominator of a fraction exceeds the numerator by 2. If 5 be added to the numerator the fraction increases by unity. The fraction is.

- a. $5/7$
- b. $1/3$
- c. $7/9$
- d. $3/5$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.9 Three persons Mr. Roy, Mr. Paul and Mr. Singh together have ₹ 51. Mr. Paul has ₹ 4 less than Mr. Roy and Mr. Singh has got ₹ 5 less than Mr. Roy. They have the money as.

- a. (₹ 20, ₹ 16, ₹ 15)
- b. (₹ 15, ₹ 20, ₹ 16)
- c. (₹ 25, ₹ 11, ₹ 15)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.10 A number consists of two digits. The digits in the ten's place is 3 times the digit in the unit's place. If 54 is subtracted from the number the digits are reversed. The number is

- a. 39
- b. 92
- c. 93
- d. 94

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.11 One student is asked to divide a half of a number by 6 and other half by 4 and then to add the two quantities. Instead of doing so the student divides the given number by 5. If the answer is 4 short of the correct answer then the number was

- a. 320
- b. 400
- c. 480
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.12 If a number of which the half is greater than $\frac{1}{5}$ th of the number by 15 then the number is

- a. 50
- b. 40
- c. 80
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Simultaneous Linear Equations in Two Unknowns

- The general form of a linear equations in two unknowns x and y is $ax + by + c = 0$ where a, b are non-zero coefficients and c is a constant.
- Two such equations $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$ form a pair of simultaneous equations in x and y .
- A value for each unknown which satisfies simultaneously both the equations will give the roots of the equations.

MATHS

CA FOUNDATION

CODE : SS12

Method of Solution

Elimination Method

Cross Multiplication Method



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12

Method of Solution

Elimination Method:

Solve $2x + 5y = 9$ and $3x - y = 5$



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

$$a_1x + b_1y + c_1 = 0$$
$$a_2x + b_2y + c_2 = 0$$

Compare the ratios	Graphical representation	Algebraic interpretation
$\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$	Intersecting lines	Exactly one solution (unique)
$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$	Coincident lines	Infinitely many solutions
$\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$	Parallel lines	No solution

MATHS

EXERCISE (C)

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Que.1) The solution of the set of equations $3x + 4y = 7$, $4x - y = 3$ is

- a. $(1, -1)$
- b. $(1, 1)$
- c. $(2, 1)$
- d. $(1, -2)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.2 The values of x and y satisfying the equations $\frac{x}{2} + \frac{y}{3} = 2$, $x + 2y = 8$ are given by the pair.

- a. (3, 2)
- b. (-2, -3)
- c. (2, 3)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.3

pair.

$\frac{x}{p} + \frac{y}{q} = 2$, $x + y = p + q$ are satisfied by the values given by the

- a. $(x = p, y = q)$
- b. $(x = q, y = p)$
- c. $(x = 1, y = 1)$
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.4 The solution for the pair of equations is given by

$$\frac{1}{16x} + \frac{1}{15y} = \frac{9}{20}, \quad \frac{1}{20x} - \frac{1}{27y} = \frac{4}{45}$$

- a. $(1/4, 1/3)$
- b. $(1/3, 1/4)$
- c. $(3, 4)$
- d. $(4, 3)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.5 Solve for x and y: $\frac{4}{x} - \frac{5}{y} = \frac{x+y}{xy} + \frac{3}{10}$ and $3xy = 10(y - x)$

- a. (5, 2)
- b. (-2, -5)
- c. (2, -5)
- d. (2, 5)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.6 The pair satisfying the equations: $x+5y = 36$, $\frac{x+y}{x-y} = \frac{5}{3}$ is given by

- a. (16, 4)
- b. (4, 16)
- c. (4, 8)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.7 Solve for x and y : $x - 3y = 0$, $x + 2y = 20$.

- a. $x = 4, y = 12$
- b. $x = 12, y = 4$
- c. $x = 5, y = 4$
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.8 The simultaneous equations $7x - 3y = 31$, $9x - 5y = 41$ have solutions given by

- a. $(-4, -1)$
- b. $(-1, 4)$
- c. $(4, -1)$
- d. $(3, 7)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.9 $1.5x + 2.4y = 1.8$, $2.5(x+1) = 7y$ have solutions as

- a. $(0.5, 0.4)$
- b. $(0.4, 0.5)$
- c. $(1/2, 2/5)$
- d. $(2, 5)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.10 The values of x and y satisfying the equation $\frac{3}{x+y} + \frac{2}{x-y} = 3$,

$\frac{2}{x+y} + \frac{3}{x-y} = 3\frac{2}{3}$ are given by

- a. (1, 2)
- b. (-1, -2)
- c. (1, 1/2)
- d. (2, 1)

MATHS

EXERCISE (D)

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Que.1) $1.5x + 3.6y = 2.1$, $2.5(x+1) = 6y$

- a. $(0.2, 0.5)$
- b. $(0.5, 0.2)$
- c. $(2, 5)$
- d. $(-2, -5)$

MATHS

CA FOUNDATION

CODE : SS12

Que.2

$$\frac{x}{5} + \frac{y}{6} + 1 = \frac{x}{6} + \frac{y}{5} = 28$$

- a. (6, 9)
- b. (9, 6)
- c. (60, 90)
- d. (90, 60)



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.3 $\frac{x}{4} = \frac{y}{3} = \frac{z}{2}; 7x + 8y + 5z = 62$

- a. (4, 3, 2)
- b. (2, 3, 4)
- c. (3, 4, 2)
- d. (4, 2, 3)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.4

$$\frac{xy}{x+y}=20, \frac{yz}{y+z}=40, \frac{zx}{z+x}=24$$

- a. (120, 60, 30)
- b. (60, 30, 120)
- c. (30, 120, 60)
- d. (30, 60, 120)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.5 $2x + 3y + 4z = 0$, $x + 2y - 5z = 0$, $10x + 16y - 6z = 0$

- a. $(0, 0, 0)$
- b. $(1, -1, 1)$
- c. $(3, 2, -1)$
- d. $(1, 0, 2)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.6) $\frac{1}{3}(x+y) + 2z = 21, 3x - \frac{1}{2}(y+z) = 65, x + \frac{1}{2}(x+y-z) = 38$

- a. (4, 9, 5)
- b. (2, 9, 5)
- c. (24, 9, 5)
- d. (5, 24, 9)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.7)

$$\frac{4}{x} - \frac{5}{y} = \frac{x+y}{xy} + \frac{3}{10} \quad 3xy = 10(y-x)$$

- a. (2, 5)
- b. (5, 2)
- c. (2, 7)
- d. (3, 4)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.8)

$$\frac{x}{0.01} + \frac{y+0.03}{0.05} = \frac{y}{0.02} + \frac{x+0.03}{0.04} = 2$$

- a. (1, 2)
- b. (0.1, 0.2)
- c. (0.01, 0.02)
- d. (0.02, 0.01)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.9)

$$\frac{xy}{y-x}=110, \quad \frac{yz}{z-y}=132, \quad \frac{zx}{z+x}=\frac{60}{11}$$

- a. (12, 11, 10)
- b. (10, 11, 12)
- c. (11, 10, 12)
- d. (12, 10, 11)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Que.10) $3x - 4y + 70z = 0$, $2x + 3y - 10z = 0$, $x + 2y + 3z = 13$

- a. $(1, 3, 7)$
- b. $(1, 7, 3)$
- c. $(2, 4, 3)$
- d. $(-10, 10, 1)$

MATHS

EXERCISE (E)

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Choose the most appropriate option (a), (b), (c) or (d).

Que.1) Monthly incomes of two persons are in the ratio 4 : 5 and their monthly expenses are in the ratio 7 : 9. If each saves ₹50 per month find their monthly incomes.

- a. (500, 400)
- b. (400, 500)
- c. (300, 600)
- d. (350, 550)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.2) Find the fraction which is equal to $\frac{1}{2}$ when both its numerator and denominator are increased by 2. It is equal to $\frac{3}{4}$ when both are increased by 12.

- a. $\frac{3}{8}$
- b. $\frac{5}{8}$
- c. $\frac{2}{8}$
- d. $\frac{2}{3}$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.3) The age of a person is twice the sum of the ages of his two sons and five years ago his age was thrice the sum of their ages. Find his present age.

- a. 60 years
- b. 52 years
- c. 51 years
- d. 50 years

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.4) A number between 10 and 100 is five times the sum of its digits. If 9 be added to it the digits are reversed find the number.

- a. 54
- b. 53
- c. 45
- d. 55

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.5) The wages of 8 men and 6 boys amount to 33. If 4 men earn ₹ 4.50 more than 5 boys determine the wages of each man and boy.

- a. (₹ 1.50, ₹ 3)
- b. (₹ 3, ₹ 1.50)
- c. (₹ 2.50, ₹ 2)
- d. (₹ 2, ₹ 2.50)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.6) A number consisting of two digits is four times the sum of its digits and if 27 be added to it the digits are reversed. The number is:

- a. 63
- b. 35
- c. 36
- d. 60

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.7) Of two numbers, $\frac{1}{5}^{\text{th}}$ of the greater is equal to $\frac{1}{3}^{\text{rd}}$ of the smaller and their sum is 16. The numbers are:

- a. (6, 10)
- b. (9, 7)
- c. (12, 4)
- d. (11, 5)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.8) y is older than x by 7 years 15 years back x's age was $\frac{3}{4}$ of y's age. Their present ages are:

- a. $(x = 36, y = 43)$
- b. $(x = 50, y = 43)$
- c. $(x = 43, y = 50)$
- d. $(x = 40, y = 47)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.9) The sum of the digits in a three digit number is 12. If the digits are reversed the number is increased by 495 but reversing only of the ten's and unit digits increases the number by 36. The number is

- a. 327
- b. 372
- c. 237
- d. 273

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.10) Two numbers are such that twice the greater number exceeds twice the smaller one by 18 and $\frac{1}{3}^{\text{rd}}$ of the smaller and $\frac{1}{5}^{\text{th}}$ of the greater number are together 21. The numbers are:

- a. (36, 45)
- b. (45, 36)
- c. (50, 41)
- d. (55, 46)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.11) The demand and supply equations for a certain commodity are $4q + 7p = 17$ and $p = \frac{q}{3} + \frac{7}{4}$ respectively where p is the market price and q is the quantity then the equilibrium price and quantity are:

- a. $2, \frac{3}{4}$
- b. $3, \frac{1}{2}$
- c. $5, \frac{3}{4}$
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Quadratic Equation

- An equation of the form $ax^2 + bx + c = 0$ where x is a variable and a, b, c are constants with $a \neq 0$ is called a quadratic equation or equation of the second degree.
- When $b = 0$ the equation is called a pure quadratic equation
- when $b \neq 0$ the equation is called an affected quadratic.

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Quadratic Equation

- **Examples:**

i) $2x^2 + 3x + 5 = 0$

ii) $x^2 - x = 0$

iii) $5x^2 - 6x - 3 = 0$

- The value of the variable say x is called the root of the equation.
- A quadratic equation has got two roots.

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

How to find out the roots of a quadratic equation:

- Factorisation Method
- Quadratic Formula

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

How to find out the roots of a quadratic equation:

$$ax^2 + bx + c = 0$$

$$\text{or } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Let one root be α and the other root be β

$$\alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

$$\beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Sum and Product of the Roots:

$$ax^2 + bx + c = 0$$

sum of roots = $-\frac{b}{a} = -\frac{\text{coefficient of } x}{\text{coefficient of } x^2}$

product of the roots = $\frac{c}{a} = \frac{\text{constant term}}{\text{coefficient of } x^2}$

MATHS

CA FOUNDATION

CODE : SS12

How To Construct A Quadratic Equation

$$x^2 - (\text{sum of the roots})x + \text{Product of the roots} = 0$$



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Nature of the roots

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- If $b^2 - 4ac = 0$ the roots are real and equal;
- If $b^2 - 4ac < 0$ then the roots are imaginary;
- If $b^2 - 4ac > 0$ then the roots are real and unequal (or distinct);
- If $b^2 - 4ac > 0$ is a perfect square the roots are real, rational and unequal (distinct);
- If $b^2 - 4ac > 0$ but not a perfect square the roots are real, irrational and unequal.


MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma



MATHS

EXERCISE (F)

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Choose the most appropriate option (a), (b), (c) or (d).

Que.1) If the roots of the equation $2x^2 + 8x - m^3 = 0$ are equal then value of m is

- a. - 3
- b. - 1
- c. 1
- d. - 2

MATHS

CA FOUNDATION

CODE : SS12

Choose the most appropriate option (a), (b), (c) or (d).

Que.2) If $2^{2x+3} - 3^2 \cdot 2^x + 1 = 0$ then values of x are

- a. 0, 1
- b. 1, 2
- c. 0, 3
- d. 0, -3



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12

Choose the most appropriate option (a), (b), (c) or (d).

Que.3)

The values of $4 + \frac{1}{4 + \frac{1}{4 + \frac{1}{4 + \dots \infty}}}$

- a. $1 \pm \sqrt{2}$
- b. $2 + \sqrt{5}$
- c. $2 \pm \sqrt{5}$
- d. None of these



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.4) If α , β be the roots of the equation $2x^2 - 4x - 3 = 0$ the value of $\alpha^2 + \beta^2$ is

- a. 5
- b. 7
- c. 3
- d. - 4

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.5) If the sum of the roots of the quadratic equation $ax^2 + bx + c = 0$ is equal to the sum of the squares of their reciprocals then $\frac{b^2}{ac} + \frac{bc}{a^2}$

is equal to

- a. 2
- b. -2
- c. 1
- d. -1

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.6) The equation $x^2 - (p+4)x + 2p + 5 = 0$ has equal roots the values of p will be.

- a. ± 1
- b. 2
- c. ± 2
- d. -2

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.7) The roots of the equation $x^2 + (2p-1)x + p^2 = 0$ are real if.

- a. $p \geq 1$
- b. $p \leq 4$
- c. $p \geq 1/4$
- d. $p \leq 1/4$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.8) If $x = m$ is one of the solutions of the equation $2x^2 + 5x - m = 0$ the possible values of m are

- a. $(0, 2)$
- b. $(0, -2)$
- c. $(0, 1)$
- d. $(1, -1)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.9) If p and q are the roots of $x^2 + 2x + 1 = 0$ then the values of $p^3 + q^3$ becomes

- a. 2
- b. -2
- c. 4
- d. - 4

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.10) If $L + M + N = 0$ and L, M, N are rationals the roots of the equation $(M + N - L)x^2 + (N + L - M)x + (L + M - N) = 0$ are

- a. Real and irrational
- b. Real and rational
- c. Imaginary and equal
- d. Real and equal

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.11) If α and β are the roots of $x^2 = x + 1$ then value of $\frac{\alpha^2}{\beta} - \frac{\beta^2}{\alpha}$ is

- a. $2\sqrt{5}$
- b. $\sqrt{5}$
- c. $3\sqrt{5}$
- d. $-2\sqrt{5}$

MATHS

Exercise F

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Choose the most appropriate option (a), (b), (c) or (d).

Que.12) If $p \neq q$ and $p^2 = 5p - 3$ and $q^2 = 5q - 3$ the equation having roots as $\frac{p}{q}$ and $\frac{q}{p}$ is

- a. $x^2 - 19x + 3 = 0$
- b. $3x^2 - 19x - 3 = 0$
- c. $3x^2 - 19x + 3 = 0$
- d. $3x^2 + 19x + 3 = 0$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.13) If one root of $5x^2 + 13x + p = 0$ be reciprocal of the other then the value of p is

- a. -5
- b. 5
- c. $1/5$
- d. $-1/5$

MATHS

EXERCISE G

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Choose the most appropriate option (a), (b), (c) or (d).

Que1. A solution of the quadratic equation $(a + b - 2c)x^2 + (2a - b - c)x + (c + a - 2b) = 0$

- a. $x = 1$
- b. $x = -1$
- c. $x = 2$
- d. $x = -2$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 2. If the root of the equation $x^2 - 8x + m = 0$ exceeds the other by 4 then the value of m is

- a. $m = 10$
- b. $m = 11$
- c. $m = 12$
- d. $m = 9$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 3. The values of x in the equation $7(x + 2p)^2 + 5p^2 = 35xp + 117p^2$ are

- a. $(4p, -3p)$
- b. $(4p, 3p)$
- c. $(-4p, 3p)$
- d. $(-4p, -3p)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 4. The solutions of the equation $\frac{6x}{x+1} + \frac{6(x+1)}{x} = 13$ are

- a. (2, 3)
- b. (3, -2)
- c. (-2, -3)
- d. (2, -3)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 5. The satisfying of x for the equation $\frac{1}{x+p+q} = \frac{1}{x} + \frac{1}{p} + \frac{1}{q}$ are

- a. (p, q)
- b. (-p, -q)
- c. (p, -p)
- d. (-p, q)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 6. The values of x for the equation $x^2 + 9x + 18 = 6 - 4x$ are

- a. (1, 12)
- b. (-1, -12)
- c. (1, -12)
- d. (-1, 12)

MATHS

CA FOUNDATION

CODE : SS12

Choose the most appropriate option (a), (b), (c) or (d).

Que 7 The values of x satisfying the equation

$$\sqrt{(2x^2+5x-2)} - \sqrt{(2x^2+5x-9)} = 1 \text{ are}$$

- a. $(2, -9/2)$
- b. $(4, -9)$
- c. $(2, 9/2)$
- d. $(-2, 9/2)$



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 8. The solution of the equation $3x^2 - 17x + 24 = 0$ are

- a. $(2, 3)$
- b. $(2, 3\frac{2}{3})$
- c. $(3, 2\frac{2}{3})$
- d. $(3, \frac{2}{3})$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.9 The equation $\frac{3(3x^2+15)}{6} + 2x^2 + 9 = \frac{2x^2+96}{7} + 6$ has got the solution as

- a. (1, 1)
- b. (1/2, -1)
- c. (1, -1)
- d. (2, -1)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 10. The equation $\left(\frac{l-m}{2}\right)x^2 - \left(\frac{l+m}{2}\right)x + m = 0$ has got two values of x to satisfy the equation given as

- a. $\left(1, \frac{2m}{l-m}\right)$
- b. $\left(1, \frac{m}{l-m}\right)$
- c. $\left(1, \frac{2l}{l-m}\right)$
- d. $\left(1, \frac{1}{l-m}\right)$

MATHS

CODE : SS12

Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 1. The sum of two numbers is 8 and the sum of their squares is 34. Taking one number as x form an equation in x and hence find the numbers. The numbers are

- a. (7, 10)
- b. (4, 4)
- c. (3, 5)
- d. (2, 6)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que.2 The difference of two positive integers is 3 and the sum of their squares is 89. Taking the smaller integer as x form a quadratic equation and solve it to find the integers. The integers are:

- a. (7, 4)
- b. (5, 8)
- c. (3, 6)
- d. (2, 5)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 3. Five times of a positive whole number is 3 less than twice the square of the number. The number is

- a. 3
- b. 4
- c. -3
- d. 2

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 4. The area of a rectangular field is 2000 sq.m and its perimeter is 180m. Form a quadratic equation by taking the length of the field as x and solve it to find the length and breadth of the field. The length and breadth are

- a. (205m, 80m)
- b. (50m, 40m)
- c. (60m, 50m)
- d. None

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 5. Two squares have sides p cm and $(p + 5)$ cms. The sum of their squares is 625 sq. cm. The sides of the squares are

- a. (10 cm, 30 cm)
- b. (12 cm, 25 cm)
- c. (15 cm, 20 cm)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 6. Divide 50 into two parts such that the sum of their reciprocals is $\frac{1}{12}$. The numbers are

- a. (24,26)
- b. (28,22)
- c. (27, 23)
- d. (20, 30)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 7. There are two consecutive numbers such that the difference of their reciprocals is $1/240$. The numbers are

- a. (15, 16)
- b. (17, 18)
- c. (13, 14)
- d. (12, 13)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 8. The hypotenuse of a right-angled triangle is 20 cm. The difference between its other two sides be 4cm. The sides are

- a. (11cm, 15cm)
- b. (12cm, 16cm)
- c. (20cm, 24cm)
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 9. The sum of two numbers is 45 and the mean proportional between them is 18. The numbers are

- a. (15, 30)
- b. (32, 13)
- c. (36, 9)
- d. (25, 20)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 10. The sides of an equilateral triangle are shortened by 12 units 13 units and 14 units respectively and a right angle triangle is formed. The side of the equilateral triangle is

- a. 17 units
- b. 16 units
- c. 15 units
- d. 18 units

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 11. A distributor of apple Juice has 5000 bottle in the store that it wishes to distribute in a month. From experience it is known that demand D (in number of bottles) is given by $D = -2000p^2 + 2000p + 17000$. The price per bottle that will result zero inventory is

- a. ₹ 3
- b. ₹ 5
- c. ₹ 2
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que12. The sum of two irrational numbers multiplied by the larger one is 70 and their difference is multiplied by the smaller one is 12; the two numbers are

- a. $3\sqrt{2}, 2\sqrt{3}$
- b. $5\sqrt{2}, 3\sqrt{5}$
- c. $2\sqrt{2}, 5\sqrt{2}$
- d. None of these

MATHS

CA FOUNDATION

CODE : SS12

Cubic Equation

$$ax^3 + bx^2 + cx + d = 0$$

where a is coefficient of x^3

where b is coefficient of x^2

where c is coefficient of x

where d is constant

$$a \neq 0$$



Shivani
Sharma

MATHS

CA FOUNDATION

CODE : SS12

Solution Of Cubic Equation

Example Solve $x^3 - 7x + 6 = 0$



Shivani
Sharma

MATHS

EXERCISE I

CODE : SS12



Shivani
Sharma

CA FOUNDATION

Choose the most appropriate option (a), (b), (c) or (d).

Que 1. The solution of the cubic equation $x^3 - 6x^2 + 11x - 6 = 0$ is given by the triplet :

- a. $(-1, 1, -2)$
- b. $(1, 2, 3)$
- c. $(-2, 2, 3)$
- d. $(0, 4, -5)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 2. The cubic equation $x^3 + 2x^2 - x - 2 = 0$ has 3 roots namely.

- a. (1, -1, 2)
- b. (-1, 1, -2)
- c. (-1, 2, -2)
- d. (1, 2, 2)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 3. x , $x - 4$, $x + 5$ are the factors of the left-hand side of the equation.

- a. $x^3 + 2x^2 - x - 2 = 0$
- b. $x^3 + x^2 - 20x = 0$
- c. $x^3 - 3x^2 - 4x + 12 = 0$
- d. $x^3 - 6x^2 + 11x - 6 = 0$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 4. The equation $3x^3 + 5x^2 = 3x + 5$ has got 3 roots and hence the factors of the left-hand side of the equation $3x^3 + 5x^2 - 3x - 5 = 0$ are

- a. $x - 1, x - 2, x - 5/3$
- b. $x - 1, x + 1, 3x + 5$
- c. $x + 1, x - 1, 3x - 5$
- d. $x - 1, x + 1, x - 2$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 5. The roots of the equation $x^3 + 7x^2 - 21x - 27 = 0$ are

- a. $(-3, -9, -1)$
- b. $(3, -9, -1)$
- c. $(3, 9, 1)$
- d. $(-3, 1, 9)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 6. The roots of $x^3 + x^2 - x - 1 = 0$ are

- a. $(-1, -1, 1)$
- b. $(1, 1, -1)$
- c. $(-1, -1, -1)$
- d. $(1, 1, 1)$

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 7. The satisfying value of $x^3 + x^2 - 20x = 0$ are

- a. (1, 4, 5)
- b. (2, 4, -5)
- c. (0, - 4, 5)
- d. (0, 4, -5)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 8. The roots of the cubic equation $x^3 - 6x^2 + 9x - 4 = 0$ are

- a. (4, 1, -1)
- b. (-4, -1, -1)
- c. (-4, -1, 1)
- d. (1, 1, 4)

MATHS

CA FOUNDATION

CODE : SS12



Shivani
Sharma

Choose the most appropriate option (a), (b), (c) or (d).

Que 9. If $4x^3 + 8x^2 - x - 2 = 0$ then value of $(2x+3)$ is given by

- a. 4, -1, 2
- b. -4, 2, 1
- c. 2, -4, -1
- d. None of these.

MATHS